



Impact of Mutants of *Trichoderma* Species in Modulating Salinity Stress of Beans (*Phaseolus vulgaris* L. CV. Khomein)

Z. Rezaloo¹, S. Shahbazi^{2*}, A. A. Aliloo¹, M. Sarajuoghi³

1. Department of Plant Production and Genetics, Faculty of Agriculture, Maragheh University

2. Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute (NSTRI), Atomic Energy Organization of Iran (AEOI), Alborz, Iran.

3. Department of Agronomy, Karaj Branch, Islamic Azad University, Alborz, Iran

ABSTRACT

Salinity in Iran's agricultural soils is a growing problem and can cause yield loss in the production of many products, including beans. One of the most biocompatible and effective solutions to this problem is bio-priming with microorganisms tolerant to salinity stress (such as *Trichoderma*). This study's salinity treatment (NaCl 100 mM) doubles the tolerance threshold of the bean plant and the effect of biopriming with *Trichoderma* isolates (wild-type and mutant) has been investigated. A mutation induction method (250 Gy) was used to increase tolerance to salinity in Iranian *Trichoderma*. Growth components, germination, allometric index, seedling tissue water, and vigor were measured at the seedling stage. Results from analysis of variance and comparison of the average showed that biological priming of beans with *Trichoderma* (wild type and mutant) fungal species had a significant advantage over control of salinity stress. However, the modulating effect of salt stress was greater in seeds inoculated with mutant species than in their wild type.

Keywords: Biopriming; *Trichoderma* sp.; Gamma Irradiation; Salinity Stress.

1. Introductions

Soil salinity is a condition of high salt concentrations in the soil. These soils' electrical conductivity (EC) equals four desi siemens [1]. This amount of soil salinity significantly reduces the yield of most agricultural products [2]. Presently, a considerable amount of agricultural land is somewhat saline, and the extent of saline land is increasing [3]. For example, one of the

reasons for the increasing salinity of agricultural land is improper and excessive irrigation [4]. With the increase in soil salinity, agricultural lands are destroyed. It is estimated that 50% of these lands will not be usable until the middle of the 47th century [5]. The adverse effects of salinity on plants are divided into three general groups, including the reduction of soil osmotic potential, the destruction of the physical structure of the soil,

*. Corresponding Author name: S. Shahbazi
E-mail address: sshabbazi@aeoi.org.ir

and the increase in the concentration of sodium (Na⁺) and chlorine (Cl⁻) ions [6].

Beans are one of the most important crops, essential for high protein content and use in the diet. This plant is an annual crop cultivated for seed production [7]. Regarding the cultivated area of legumes, beans are the first in Iran [8]. Beans are susceptible to salinity and tolerate soil salinity less than 2 Deci siemens/meter [9].

Many studies showed that increasing salinity in the hydroponic cultivation of beans reduced root dry weight, number of roots, total root length, average root diameter, and total root volume [10]. Also, salinity has adverse effects on bean germination; so, increasing salinity stress decreases the germination rate, germination percentage, and root length of this plant [11]. In this regard, from the results of investigating the effects of salinity stress on beans in hydroponic cultivation, it was found that with increasing salinity, fresh stem weight, root fresh and dry weight, and root length of this plant decreased [12].

Biopriming seeds can increase growth quickly, plant nutrient absorption efficiency, and relative tolerance to harsh environmental conditions such as salinity [13]. Seed biopriming causes an increase in biological and physiological activities in the primed seed and the resulting plant. This increase can be observed in the way of germination, early seedling establishment, early maturity, and quantitative and qualitative increase [14]. Using fungal biological fertilizers in seed inoculation is the most crucial method of using these fertilizers [15]. *Trichoderma* is one of the most important members of the rhizosphere microorganism community, and the positive effect of its inoculation on plant growth has been proven [16]. By penetrating the root and establishing an endophytic relationship with the plant root, this

fungus changes the gene expression profile of the plant. This changes the plant's metabolism of the plant [17, 18, 19]. There are reports of increased plant growth due to contact with *Trichoderma* [20, 21, 22]. When exposed to physiological, biotic, and abiotic stresses, the seed responds positively to *Trichoderma* treatment. Based on the study that was done on the expression of the most desirable and appropriate biological pretreatment (biopriming) in comparison with the control (no inoculation) when faced with salinity conditions in different concentrations, the salinity of agricultural soils and many reports indicate the improvement of germination behavior and indicators related to It includes average germination time, seed germ, root length, shoot length, germination rate and initial establishment in primed seeds [23, 24, 25].

Studies show the resistance of different plants treated with *Trichoderma* species to salinity stress. Considering the nutritional importance of beans, in this research, wild and mutant species of *Trichoderma* [26] were investigated. This was to study the effect of *Trichoderma* on increasing the resistance of bean plants to salt stress. Investigating the possibility of increasing salinity tolerance in bean plant seedlings by using mutant *Trichoderma* isolates which have a higher salt tolerance than their wild parent isolate has also been the secondary aim of this research.

2. Material and Methods

Nuclear Agriculture Research School-Nuclear Science and Technology Institute (NSTRI) received five *Trichoderma* isolates (Table 1) for molecular identification using ITS-rDNA gene sequencing. *Trichoderma* isolate was irradiated in the Nuclear Agriculture Research School (NSTRI, AEIO) using a cobalt-60 γ - irradiator at a dose rate of 0.018 Gy/sec following isolation and identification. The dosimetry procedure was

completed using the Fickereference standard dosimetry system (Gamma-cell Issledovatel, PX-30). To determine a suitable dose, the specified dose range was 0, 50, 150, 200, 250, 300, 350, 400, and 450 Gy. After dose optimization, irradiation was performed at an optimized dose of 250 Gy (three replicates) on spore suspension (10^6 spores/ml) of *Trichoderma*. Serial diluted (10^{-6} , 10^{-5} , 10^{-4} , 10^{-3} , 10^{-2} , 10^{-1}) spore suspensions were prepared after irradiation, and 1 ml of each diluted sample was cultured in the PDA medium and incubated at 28 °C for 72 hours [13]. To ensure the stability of the mutants, each mutant was sub-cultured seven times, and morphological characteristics were recorded for all mutants. The mutation-optimized dose was based on a spore germination inhibitory effect of approximately 40-50 % on a Water Agar (WA) medium [26]. The Seed and Plant Improvement Institute (SPII) of Iran provided the Khomein variety of pinto beans (*Phaseolus vulgaris*). Maragheh University conducted the factorial experiment in 2020 in three replications using a completely random design. The description of the treatments is given in Table 1. In comparison to the control, salt stress treatment was accomplished by irrigating water with sodium

chloride 100 mM or 9.1 ds/m. The abbreviations "S" and "M" refer to mutant and salt stress, respectively, and the results were compared to the controls. In plants, the effect of biopriming with *Trichoderma* isolates (wild type and mutant) has been investigated to double the tolerance threshold (5 ds/m) of bean plants [27].

Seeds were superficially disinfected in ethanol 70% for 1 minute and sodium hypochlorite 2% for 40 seconds, and finally, washed 3-4 times with sterile distilled water. Then the seeds were inoculated in spore suspension with a population of 1×10^8 for 20 minutes on a shaker. Bean seeds were planted in 3-liter pots containing washed perlite, and five seeds were planted in each pot. The pots were kept in the growth chamber under controlled conditions under a lighting period of 8.16 hours (dark/light), a temperature period of 18.28 degrees Celsius (night/day), and relative humidity of 50-60%. It was irrigated with distilled water (irrigation was twice a week) and Hoagland's nutrient solution (Hoagland and Arnon, 1950) [28] with an approximate pH of 7.5 ± 1 . The pots were subjected to salt treatment at two concentration levels of zero (control) and 100 mM sodium chloride.

Table 1. Treatments used in this research, *Trichoderma* sp. with NCBI code.

NCBI Code	NSTRI Code	Treatments	
MW718882	NAS107	<i>T. harzianum</i>	<i>T. harzianum</i> * S
MW719563	NAS114	<i>T. lixii</i>	<i>T. lixii</i> * S
MW719590	NAS108	<i>T. ghanens</i>	<i>T. ghanens</i> * S
MW719876	NAS115	<i>T. virens</i>	<i>T. virens</i> * S
MW719255	NAS112	<i>T. atroviride</i>	<i>T. atroviride</i> * S
MW719591	NAS107 M8	<i>T. harzianum</i> mutant	<i>T. harzianum</i> mutant * S
MW719564	NAS114 M17	<i>T. lixii</i> mutant	<i>T. lixii</i> mutant* S
ON545796)	NAS108 M1	<i>T. ghanens</i> mutant	<i>T. ghanens</i> mutant* S
MW719878	NAS115 M17	<i>T. virens</i> mutant	<i>T. virens</i> mutant * S
MW719257	NAS112M2	<i>T. atroviride</i> mutant	<i>T. atroviride</i> mutant* S
		Control	Salt stress (S)

At the end of the experiment, the factors of length, fresh and dry weight of shoots and roots, the relative percentage of leaf water, and emergence were investigated.

The method of Cherki et al. [26, 29] was used to measure the percentage of water in fresh leaves. For this purpose, in each replication, a disc with a diameter of one centimeter was prepared from the leaf and then weighed with a scale with an accuracy of 0.0001 gram (FW). The leaf discs were placed in distilled water for 4 hours, and their weight was determined again (TW). These disks were dried in an oven at 70 degrees Celsius, and their dry weight was obtained (DW). The following relationship was used for seedling tissue water percentage;

$$RWC = \frac{FW-DW}{TW-DW} \times 100 \quad (1)$$

The following equation was used to calculate the vigor index [26];

(2)

Vigor Index

$$= \frac{\text{Germination Percentage} \times \text{Seedling dry Weight}}{100}$$

Also, the allometric coefficient was calculated using the following equation [28];

$$\text{Allometric coefficient} = \frac{\text{Stem Length}}{\text{Root Length}} \quad (3)$$

The obtained data were analyzed through the SPSS version 9 statistical program and, Duncan's multi-range test was used to compare the mean at a 5% probability level. Graphs were also drawn using Excel software.

3. Results and Discussion

The analysis of variance showed that the length, wet and dry weight of shoots and roots under the influence of salt stress and bioprimering and the mutual effects of stress and bioprimering showed a statistically significant difference at the probability level of 1% (Table 2). Also, the allometric index, seedling tissue water percentage, and seed vigor showed statistically significant differences under the effect of salinity stress and bioprimering and the mutual effects of stress and bioprimering at the probability level of 1% (Table 2). Germination rate was significant at the probability level of 1% for salinity stress and bioprimering with *Trichoderma* species and was not significant for the interaction effect of salinity stress and bioprimering (Table 2).

Table 2. Analysis of variance (mean square) of bean characteristics under salt stress and bioprimering treatments.

Source of Variation	df	percentage of Emergence	Length		Fresh Weight		Dry Weight		Allometric Index	Seedling Tissue Water Percentage	Seed Vigor
			Root	Stem	Root	Stem	Root	Stem			
bio-primering	10	321.343**	20.83**	234.59**	5.78**	35.34**	0.84**	0.06**	0.08**	19.64**	84356.4635**
Salt stress	1	6060.325**	30.93**	27.27**	2.43**	2.64**	0.25**	0.005**	0.14**	6.47**	54064.8336**
bio-primering*Salt stress	10	55.36 ^{ns}	0.14**	0.26**	0.036**	0.007**	0.06**	0.0002**	0.08**	0.52**	435.3597**
Error	10	39.58	0.05	0.05	0.009	0.00003	0.0007	0.00003	0.003	0.03	345.6735
CV.		10.04	2.51	1.06	3.83	1.89	6.94	5.51	12.49	0.12	9.73

^{ns}, ** and *: non-significance, respectively, significant at the probability level of 1 and 5 percent.

Germination rate

The average comparison between the treatments showed that the 100% germination rate of seeds was observed in *T. atroviride* mutant, *T. ghanens*, *T. ghanens* mutant, *T. harzianum*, and *T. harzianum* mutant without stress. The seeds treated with *T. atroviride*, and *T. virens*, had no significant difference (99% germination rate). The lowest germination was in the saline stress treatment (69%) which caused a 24.2% decrease in germination compared to the control (91%) (Fig. 1).

In the interaction between salinity and *Trichoderma*, *T. virens* mutant (97%) had the highest salinity stress modulating effect (*T. virens* mutant*S 89%). The effect of salinity stress in *T. virens* mutant*S was 8% less than in *T. virens* mutant (Fig. 1).

The *T. harzianum* *S had the highest salinity stress modulating effect (88%) and was able to reduce the stress effect by 21% compared to S (NaCl 100 mM) (69%) (Fig. 1). Salinity stress in *T. harzianum**S treatment reduced germination by 12% compared to *T. harzianum* (100%) (Fig. 1).

All *Trichoderma* species facilitate the growth and germination of plants in saline soils [30]. Different species of *Trichoderma* fungi affect plant growth and are called growth promoters [31]. Research showed that the inoculation of bean plant seeds with *Trichoderma* fungus could increase the germination rate under salt stress compared to the control [32]. The use of *Trichoderma* in bean plants increased the percentage and speed of germination compared to the control [33]. Salinity reduces the germination rate by reducing water potential and the effect of absorbed ions on active enzymes and hormones inside the seed [34]. Growth-promoting fungi increase the germination rate by producing indole acetic acid in the growth medium [35]. Ions

in irrigation water during germination can stimulate, inhibit or neutralize germination [36]. The lower value of the average germination time means that the seeds have germinated in a shorter period. The seed's water absorption is disrupted or slow in conditions of stress. In such a case, the metabolic activities of germination are carried out slowly inside the seed, and as a result, the time required for germination increases [37]. It has been reported that the use of biofertilizers and their combination in basil medicinal plants improved germination indicators such as percentage and average duration of germination [38].




Fig. 1. Emergence percentage of bean seeds in salinity stress treatments, bioprimering, and interaction effect of salinity and bioprimering.

Seedlings treated with *T. lixii* and *T. harzianum* had the longest roots, 17.83 cm and 17.84 cm, respectively. The stem length in the *T. virens* and *T. harzianum* treatments was 30.83 and 29.62 cm, in turn, which had the longest stem length among the treatments. In S treatment, the length of the stem and root was 15.87 cm and 5.73 cm, respectively, and had the highest inhibitory effect on the length of the stem and root (Fig. 2). Among the treatments under stress, *T. virens* *S had a root length of 12.26 cm and compared to *T. virens* with a root length of 16.73 cm, it decreased by 26.72%. It was reported that mutant S (6.83 cm) had the lowest root length, however, this isolate was able to control the stress by 16.11% more than mutant S

(Fig 2). The results of various experiments show that both the length of the root and the stem decreased as a result of salt stress, but the reduction ratio of the length of the stem is more than the length of the root [39]. Many reports indicate the improvement of germination rate, vigor, root length, stem length, and initial establishment in bio-primed seeds with *Trichoderma* species [40]. The studies of Shafi et al. [41] that investigated the effect of salinity stress (NaCl) on wheat root morphology showed that NaCl reduced root dry weight, root number, total root length, average root diameter, and total root volume.




Fig 2. Length of bean seedling in salinity stress treatments, biopriming, and interaction effect of salinity and biopriming.

Fresh Weight

The fresh weight of the root in the seedlings treated with *T. lixii* was 18.74 g, compared to the control (14.03 g), 25.14%, and in comparison to S (7.34 g), 60.84%. *T. virens* mutant *S (13.45 g) had the greatest modulating effect on root fresh weight. This mutant had an increase of 1.86% compared to its wild type (*T. virens**S 13.2 g). S (7.34 g) had the lowest root fresh weight and *T. ghanens* (12.96 g) had the lowest root fresh weight among the wild type. Of course, the effect of gamma radiation in the *T. ghanens* mutant (13.3 g) increased the root fresh weight by 2.94% as opposed to the wild type

of *T. ghanens*. The lowest stress-modulating effect was observed in *T. ghanens* *S (8.63 g) and gamma-ray increased the root fresh weight by 5.09% in *T. ghanens* *S (9.07 g).

There was a significant increase in stem fresh weight (51.73 cm) in *T. atroviride* mutant seedlings treated with *Trichoderma* without salinity stress. The lowest stem fresh weight among the treatments was related to S (22.03). Among the treatments under salinity stress, *T. lixii* *S with 41.71 grams of stem fresh weight had the greatest increase and modulating effect of salinity stress and was able to neutralize the effect of salinity by 89.33% compared to S. The lowest stem fresh weight (28.89) was related to *T. atroviride* *S, and the effect of gamma rays in this *T. atroviride* mutant *S species could increase the neutralizing effect of salinity stress by 30.52% (Fig. 3). Sánchez-Montesinos et al. [42] reported that the application of *Trichoderma* fungus leads to an increase in the growth parameters of legumes under salinity stress. Many published studies investigated the effect of some isolates of *Trichoderma* fungus on beans and showed that this fungus improved the growth characteristics of beans under greenhouse conditions [43]. The researchers investigated the response to irrigation water salinity at different stages of root growth and found that the highest root yield and crop yield belonged to the control treatment that received the lowest irrigation water salinity increasing water salinity decreased root yield and crop performance [44]. Also, Zhang et al. in several studies [45] investigated the effects of sodium chloride salinity stress on the yield, quantity, and quality of fruit in a part of the root system. It showed that by applying salinity to the whole root system, the fruit's fresh and dry weight, the fruit's length, and the fruit's yield decreased significantly.




Fig. 3. Fresh Weight of bean seedling in salinity stress treatments, biopriming, and interaction effect of salinity and biopriming.

Dry Weight

The highest dry weight of root (2.55 g) of *T. virens* and stem in *T. virens*, *T. virens* mutant, *T. lixii*, *T. atroviride* mutant, and control (respectively: 7.04, 6.88, 6.77, 6.77, 6.79 g) was observed. The lowest dry weight of the root and stem in S was 1.00 and 3.00. Among the non-stressed treatments, the dry weight of the wild *T. virens* species was the highest, and the ratio of the control was 13.33% and 3.68% higher. *T. virens* mutant roots and stems had the highest dry weight. Among the treatments under the stress of the wild type, *T. harzianum* *S (1.84 g) had the highest salinity modulating effect, which increased root dry weight by 84% compared to S (1.00 g). The dry weight of the stem in *T. harzianum* *S (5.17 g) was higher than other wild species and it was 72.33% higher than S (Graph. 4). *T. atroviride* mutant *S had the highest stem dry weight (5.68 g) among the mutant species. The effect of gamma radiation in this species increased the dry weight of the stem by 45.64% compared to the wild type under the same conditions. The modulating effect of *T. atroviride* mutant *S was 89.33% higher than S (Graph. 4). Rezaloo et al. [20, 22, 26] reported the increasing effect of wet and dry root weight and seedling length on bean, soybean, rapeseed, lettuce, sugar beet, wheat, corn,

tomato, and cucumber seeds coated with *Trichoderma* species (wild type and mutant). The studies by Tabatabaie et al. [46] who studied the effect of uneven distribution of salt on the root and on the ratio of the product to the amount of water consumed in the hydroponic cultivation of tomato plants showed that in the hydroponic system with mobile branched roots in which the replacement of the root branches between the nutrient solution and salt water is applied with an interval of 3 days, a significant decrease in the growth of root and aerial organs was observed.




Fig. 4. Dry Weight of bean seedling in salinity stress treatments, biopriming, and interaction effect of salinity and biopriming (abbreviations: m: mutant, s: salt stress, Th: *T. harzianum*, Tl, *T. lixii*, Tg: *T. ghanens*, Tv: *T. virens*, Ta: *T. atroviride*).

Allometric Index

The allometric coefficient of *T. atroviride* *S, *T. atroviride*, and *T. harzianum* mutant *S was higher than in other treatments. *T. ghanens* had the lowest allometric coefficient. (Table 3). In the conditions of not using *Trichoderma*, the allometric coefficient increased with the increase in salinity. However, the use of *Trichoderma* in saline conditions caused a decrease in the allometric coefficient, which means an increase in the length of the rhizome compared to the stem. This trait may be contributing to resistance to salinity or subsequent drought. The use of *Trichoderma* in inoculation with bean seeds increased root length,

stem length, and dry matter. It also showed characteristics such as organic phosphorus mineralization, auxin production, ACC deaminase activity, ammonia production, and siderophore production [47].

Seedling Tissue Water Percentage

The average water potential of the plant tissue between treatments varied from 18.92% to 44.55%. The *T. virens* mutant *S had the highest seed tissue water percentage, 135.46% higher than the control at the same level. Salinity stress causes a kind of physiological dryness by reducing the osmotic potential of soil water. Such physiological dryness can cause disturbances in photosynthesis and plant life [48]. The decrease in the relative water percentage of leaves is the general response of rice plants exposed to osmotic stress. This is a very reliable indicator of the water status of the plant [49]. In glycophytes, high salinity reduces water stress and consequently reduces growth, one of the most important effects of high salinity [50]. By using Trichoderma fungi, the relative water content index was improved (Table 3). Yeşilyurt et al [51] reported that the biopriming of the seed with Trichoderma increased the relative water content under salinity stress conditions.

Vigor Index

A comparison of the average between the treatments showed that the highest amount of root index was related to the treatment of seeds with *T. ghanens* mutant without salt stress, which increased by 106.92% compared to the control. S had the lowest vigor index, which decreased by 7.93% compared to the control (Table 3). The study of seed treatment with growth-promoting fungi on the germination indices of soybean plants under salt stress by Bakhshandeh et al. [52] showed

that seed inoculation with Trichoderma had a positive effect on the germination indices and seed germination. Since Trichoderma can produce siderophore and indole acetic acid, it also has ACC deaminase activity and increases the growth of rice plants [53]. *T. harzianum* was isolated from the rhizosphere of the potato plants. It showed the ability to produce a siderophore, dissolve phosphate, and produce indole acetic acid, hydrogen cyanide, ammonia, and exopolysaccharide. This fungus increases the plant's growth activities even in soil contaminated with fungicides [54].

Table 3. Comparison of the average growth components of bean seeds under salinity stress and biopriming treatments and the mutual effect of salinity stress and biopriming.

treatment	Alometric Index	Seedling Tissue Water Percentage	Vigor Index
Control	0.31 cd	42.92 b	441.57 ijk
<i>T. atroviride</i>	0.40 a	27.42 kl	881.82 b
<i>T. atroviride</i> *NaCl	0.41 a	24.63 mn	858.99 bc
<i>T. atroviride</i> m	0.37 b	42.82 b	598.62 g
<i>T. atroviride</i> m*NaCl	0.31 cd	35.90 de	641.46 f
<i>T. ghanens</i>	0.24 g	31.86 i	888.96 b
<i>T. ghanens</i> *NaCl	0.37 b	31.68 i	470.55 ij
<i>T. ghanens</i> m	0.33 c	27.06 l	913.07 a
<i>T. ghanens</i> m*NaCl	0.37 b	35.06 f	738.33 d
<i>T. harzianum</i>	0.33 c	41.96 c	867.40 bc
<i>T. harzianum</i> *NaCl	0.34 c	32.69 g	879.97 b
<i>T. harzianum</i> m	0.37 b	32.45 gh	694.14 e
<i>T. harzianum</i> m*NaCl	0.41 a	27.30 kl	429.38 jk
<i>T. lixii</i>	0.26 f	42.85 b	494.82 hi
<i>T. lixii</i> *NaCl	0.33 c	30.99 j	576.79 gh
<i>T. lixii</i> m	0.34 c	41.38 cd	538.66 ghi
<i>T. lixii</i> m*NaCl	0.40 a	27.913 k	739.12 d
<i>T. virens</i>	0.37 b	30.83 j	276.09 m
<i>T. virens</i> *NaCl	0.29 e	24.84 m	837.93 c
<i>T. virens</i> m	0.29 e	43.54 ab	508.17 hi
<i>T. virens</i> m*NaCl	0.29 e	44.55 a	496.17 hi
NaCl (100 mM)	0.33 c	18.92 o	409.41 kl

In each column, the averages that have at least one letter in common have no significant difference based on Duncan's test at the 5% probability level.



Fig. 1. Comparison of the mutual effect of Th*S (*Trichoderma harzianum*), Th m*S (*Trichoderma harzianum* mutant), and S (salinity stress) on bean plant.



Fig. 2. Comparison of bean plants treated with *Trichoderma*. The control bean plant is shorter and weighs less compared to Tg (*Trichoderma ghanens*). The bean plant treated with Tg m (*Trichoderma ghanens* Mutant) has a higher fresh weight and more freshness.

4. Conclusions

The *Trichoderma* fungus has been effective in improving the growth of bean plants [22]. In addition, it should not be overlooked that plant seed coating with biological agents such as *Trichoderma*, controls a wide range of fungal and bacterial agents [26]. This can be proposed as a biotechnological approach to enhance growth and resistance to biotic and abiotic stresses. With the *Trichoderma* fungus, the plant seems to be able to deal better with salt stress. As seen in this study, it can be shown that the induction of gamma-ray mutation and the selection of suitable mutants can reduce the initial reactions of some plants to the presence of *Trichoderma* in the ecological root niche and increase the positive effects of this fungus; so, it is recommended as a promising technique. Although the physiological changes of the plant depend on the method of inoculation and

different isolates of *Trichoderma* used, it should not be forgotten that before recommending this biological agent, the interactions of different species of this fungus should be investigated independently and then by each plant (and even any varietal variety). It is suggested to investigate the physiological mechanisms of creating this tolerance to salinity in the plant treated with *Trichoderma*, the study of mutated genes in *Trichoderma* mutants that have a higher tolerance to salinity, and the interaction studies of the plant under salt stress and treated with this biological agent in the future studies will be done.

5. Acknowledgments

This article is a part of the Ph.D. thesis of the first author, and the authors are grateful to the Nuclear Science and Technology Research Institute (NSTRI), Atomic Energy Organization of Iran (AEOI) for their support of the implementation of this project.

References

1. Bughici T, Skaggs TH, Corwin DL, Scudiero E. Ensemble HYDRUS-2D modeling to improve apparent electrical conductivity sensing of soil salinity under drip irrigation. [Agricultural Water Management](#). 2022 Oct 1;272:107813.
2. Slamini M, Sbaa M, Arabi M, Darmous A. Review on Partial Root-zone Drying irrigation: Impact on crop yield, soil and water pollution. [Agricultural Water Management](#). 2022 Sep 1;271:107807.
3. Aboelsoud HM, AbdelRahman MA, Kheir AM, Eid MS, Ammar KA, Khalifa TH, Scopa A. Quantitative estimation of saline-soil amelioration using remote-sensing indices in arid land for better management. [Land](#). 2022 Jul 8;11(7):1041.
4. Hossain MS. Present scenario of global salt affected soils, its management and importance of salinity research. [Int. Res. J. Biol. Sci](#). 2019 Jan;1(1):1-3.
5. Brambati A. Coastal zone problems and management: a brief review. [Chemistry and Ecology](#). 2004 Jun 1;20(sup1):155-66.

6. El-Mageed TA, Mekdad AA, Rady MO, Abdelbaky AS, Saady HS, Shaaban A. Physio-biochemical and agronomic changes of two sugar beet cultivars grown in saline soil as influenced by potassium fertilizer. [Journal of Soil Science and Plant Nutrition](#). 2022 Sep;22(3):3636-54.
7. Ma J, Khan N, Gong J, Hao X, Cheng X, Chen X, Chang J, Zhang H. From an introduced pulse variety to the principal local agricultural industry: A case study of red kidney beans in Kelan, China. [Agronomy](#). 2022 Jul 21;12(7):1717.
8. FAO. FAO Production Year Book (58), Food and Agricultural Organization of the United Nations, Rome, Italy. 2021
9. Alsuvaïd M, Demir Y, Kiremit MS, Arslan H. Interaction effect of water magnetization and water salinity on yield, water productivity and morpho-physiological of balkız bean (*Phaseolus vulgaris*). [Gesunde Pflanzen](#). 2022 Jun 1;74(3):00606.
10. Rafat A, Mostafa D. Differential responses of common bean, (*Phaseolus vulgaris* L.) to the interactive effects of ascorbic acid and *Trichoderma harzianum* under salinity stress. [Egyptian Academic Journal of Biological Sciences](#), H. Botany. 2022 Jul 22;13(2):11-29.
11. Nasrallah AK, Kheder AA, Kord MA, Fouad AS, El-Mogy MM, Atia MA. Mitigation of salinity stress effects on Broad bean productivity using calcium phosphate nanoparticles application. [Horticulturae](#). 2022 Jan 14;8(1):75.
12. Santos MG, Ribeiro RV, Machado EC, Pimentel C. Photosynthetic parameters and leaf water potential of five common bean genotypes under mild water deficit. [Biologia Plantarum](#). 2009 Jun;53:229-36.
13. Xi L, Zhang M, Zhang L, Lew TT, Lam YM. Novel materials for urban farming. [Advanced Materials](#). 2022 Jun;34(25):2105009.
14. Acharya P, Jayaprakash GK, Crosby KM, Jifon JL, Patil BS. Nanoparticle-mediated seed priming improves germination, growth, yield, and quality of watermelons (*Citrullus lanatus*) at multi-locations in Texas. [Scientific reports](#). 2020 Mar 19;10(1):5037.
15. Chen K, Scheper J, Fijen TP, Kleijn D. Potential tradeoffs between effects of arbuscular mycorrhizal fungi inoculation, soil organic matter content and fertilizer application in raspberry production. [Plos one](#). 2022 Jul 18;17(7):e0269751.
16. Shahriar SA, Islam MN, Chun CN, Kaur P, Rahim MA, Islam MM, Uddain J, Siddiquee S. Microbial metabolomics interaction and ecological challenges of *Trichoderma* species as biocontrol inoculant in crop rhizosphere. [Agronomy](#). 2022 Apr 8;12(4):900.
17. Doni F, Fathurrahman F, Mispan MS, Suhaimi NS, Yusoff WM, Uphoff N. Transcriptomic profiling of rice seedlings inoculated with the symbiotic fungus *Trichoderma asperellum* SL2. [Journal of Plant Growth Regulation](#). 2019 Dec;38:1507-15.
18. Nogueira-Lopez G, Greenwood DR, Middleditch M, Winefield C, Eaton C, Steyaert JM, Mendoza-Mendoza A. The apoplasmic secretome of *Trichoderma virens* during interaction with maize roots shows an inhibition of plant defence and scavenging oxidative stress secreted proteins. [Frontiers in plant science](#). 2018 Apr 5;9:409.
19. Harman G, Khadka R, Doni F, Uphoff N. Benefits to plant health and productivity from enhancing plant microbial symbionts. [Frontiers in Plant Science](#). 2021 Apr 12;11:610065.
20. Rezaloo Z, Shahbazi S, Askari H. Biopriming with *Trichoderma* on Germination and vegetative characteristics of sweet corn, sugar beet and wheat. [Iranian Journal of Seed Science and Technology](#). 2020 Jan 21;8(2):199-210.
21. Rezaloo Z, Tohidloo G, Shahbazi S. Study of yield, vegetative traits and seed germination of pinto bean produced from biopriming with *Trichoderma*. [Journal of Agronomy and Plant Breeding](#). 2018; 14(2): 39-47.
22. Rezaloo Z, Tohidloo G, Shahbazi S, Askari H. Effects of Biopriming with *Trichoderma* Fungi on Germination and Some Vegetative characteristics of Soybean and Canola. [Journal of Agronomy and Plant Breeding](#). 2018; 14(1): 39-48.
23. Sen A, Johnson R, Puthur JT. Seed priming: A cost-effective strategy to impart abiotic stress tolerance. [Plant Performance Under Environmental Stress: Hormones, Biostimulants and Sustainable Plant Growth Management](#). 2021:459-80.
24. Kumar V, Singh G, Dwivedi SK. Arsenic and Nickel Stress Response, Their Bioremediation Potential and Mechanism of *Trichoderma* lixii Isolated from Electroplating Wastewater. [Geomicrobiology Journal](#). 2022 May 2;39(3-5):186-98.
25. Ogbe AA, Finnie JF, Van Staden J. The role of endophytes in secondary metabolites accumulation in medicinal plants under abiotic stress. [South African Journal of Botany](#). 2020 Nov 1;134:126-34.

26. Ghasemi S, Safaie N, Shahbazi S, Shams-Bakhsh M, Askari H. Enhancement of Lytic Enzymes Activity and Antagonistic Traits of *Trichoderma harzianum* Using γ -Radiation Induced Mutation. [Journal of Agricultural Science and Technology](#). 2019 Jul 10;21(4):1035-48.
27. Parande S, Zamani G, Sayyari M, Ghaderi M. Effects of silicon on the physiological, quality and quantity characteristics of common bean (*Phaseolus vulgaris*) under salinity stress. [Iranian Journal Pulses Research](#). 2014 Oct 23;5(2):57-70.
28. Hoagland DR, Arnon DI. The water-culture method for growing plants without soil. [Circular](#). California agricultural experiment station. 1950;347(2nd edit).
29. Ghoulam C, Foursy A, Fares K. Effects of salt stress on growth, inorganic ions and proline accumulation in relation to osmotic adjustment in five sugar beet cultivars. [Environmental and experimental Botany](#). 2002 Jan 1;47(1):39-50.
30. Scott SJ, Jones RA, Williams W. Review of data analysis methods for seed germination 1. [Crop science](#). 1984 Nov;24(6):1192-9.
31. Cabral-Miramontes JP, Olmedo-Monfil V, Lara-Banda M, Zúñiga-Romo ER, Aréchiga-Carvajal ET. Promotion of plant growth in arid zones by selected *Trichoderma* spp. strains with adaptation plasticity to alkaline pH. [Biology](#). 2022 Aug 12;11(8):1206.
32. López-Bucio J, Esparza-Reynoso S, Pelagio-Flores R. Nitrogen availability determines plant growth promotion and the induction of root branching by the probiotic fungus *Trichoderma atroviride* in *Arabidopsis* seedlings. [Archives of Microbiology](#). 2022 Jul;204(7):380.
33. Álvarez-García S, Rodríguez-González Á, Zanfano L, Gutiérrez S, Casquero PA. Volatile-mediated interactions between *Trichoderma harzianum* and *Acanthoscelides obtectus*: A novel in vitro methodology to evaluate the impact of microbial volatile compounds on dry grain storage pests. [Biological Control](#). 2022 Jun 1;169:104868.
34. Prisco JT, O'leary JW. Osmotic and "toxic" effects of salinity on germination of *Phaseolus vulgaris* L. seeds. [Turrialba](#). 1970;20:177-84.
35. Kaur T, Manhas RK. Evaluation of ACC deaminase and indole acetic acid production by *Streptomyces hydrogenans* DH16 and its effect on plant growth promotion. [Biocatalysis and Agricultural Biotechnology](#). 2022 Jul 1;42:102321.
36. Soliman MH, Abdulmajeed AM, Alhaithloul H, Alharbi BM, El-Esawi MA, Hasanuzzaman M, Elkelish A. Saponin biopriming positively stimulates antioxidants defense, osmolytes metabolism and ionic status to confer salt stress tolerance in soybean. [Acta Physiologiae Plantarum](#). 2020 Jul;42:1-3.
37. Lechowska K, Kubala S, Wojtyła Ł, Nowaczyk G, Quinet M, Lutts S, Garnczarska M. New insight on water status in germinating *Brassica napus* seeds in relation to priming-improved germination. [International Journal of Molecular Sciences](#). 2019 Jan 28;20(3):540.
38. Barna D, Kisvarga S, Kovács S, Csátári G, Tóth IO, Fári MG, Alshaal T, Bákonyi N. Raw and fermented alfalfa brown juice induces changes in the germination and development of french marigold (*Tagetes patula* L.) plants. [Plants](#). 2021 May 27;10(6):1076.
39. Hu DD, Dong S, Zhang J, Zhao B, Ren B, Liu P. Endogenous hormones improve the salt tolerance of maize (*Zea mays* L.) by inducing root architecture and ion balance optimizations. [Journal of Agronomy and Crop Science](#). 2022 Oct;208(5):662-74.
40. Ben-Jabeur M, Kthiri Z, Djébali N, Karmous C, Hamada W. A case study of seed biopriming and chemical priming: seed coating with two types of bioactive compounds improves the physiological state of germinating seeds in durum wheat. [Cereal Research Communications](#). 2023 Mar;51(1):125-33.
41. Shafi M, Zhang G, Bakht J, Khan MA, Islam UE, Khan MD, Raziuddin GZ. Effect of cadmium and salinity stresses on root morphology of wheat. [Pak. J. Bot.](#) 2010 Aug 1;42(4):2747-54.
42. Sánchez-Montesinos B, Diáñez F, Moreno-Gavira A, Gea FJ, Santos M. Plant growth promotion and biocontrol of *Pythium ultimum* by saline tolerant *Trichoderma* isolates under salinity stress. [International Journal of Environmental Research and Public Health](#). 2019 Jun;16(11):2053.
43. Shaban WI, El-Bramawy MA. Impact of dual inoculation with *Rhizobium* and *Trichoderma* on damping off, root rot diseases and plant growth parameters of some legumes field crop under greenhouse conditions. [Int. Res. J. Agric. Sci. Soil Sci.](#) 2011;1(3):98-108.
44. Sahin U, Ekinici M, Ors S, Turan M, Yildiz S, Yildirim E. Effects of individual and combined

- effects of salinity and drought on physiological, nutritional and biochemical properties of cabbage (*Brassica oleracea* var. *capitata*). [Scientia Horticulturae](#). 2018 Oct 20;240:196-204.
45. Zhang P, Senge M, Dai Y. Effects of salinity stress on growth, yield, fruit quality and water use efficiency of tomato under hydroponics system. [Reviews in Agricultural Science](#). 2016;4:46-55.
46. Tabatabaie SJ, Gregory PJ, Hadley P. Uneven distribution of nutrients in the root zone affects the incidence of blossom end rot and concentration of calcium and potassium in fruits of tomato. [Plant and soil](#). 2004 Jan;258:169-78.
47. Hoseini A, Salehi A, Sayyed RZ, Balouchi H, Moradi A, Piri R, Fazeli-Nasab B, Poczai P, Ansari MJ, Obaid SA, Datta R. Efficacy of biological agents and fillers seed coating in improving drought stress in anise. [Frontiers in plant science](#). 2022 Jul 22;13:955512.
48. Chourasia KN, Lal MK, Tiwari RK, Dev D, Kardile HB, Patil VU, Kumar A, Vanishree G, Kumar D, Bhardwaj V, Meena JK. Salinity stress in potato: Understanding physiological, biochemical and molecular responses. [Life](#). 2021 Jun 10;11(6):545.
49. Swapna S, Shylaraj KS. Screening for osmotic stress responses in rice varieties under drought condition. [Rice science](#). 2017 Sep 1;24(5):253-63.
50. Safdar H, Amin A, Shafiq Y, Ali A, Yasin R, Shoukat A, Hussan MU, Sarwar MI. A review: Impact of salinity on plant growth. [Nat. Sci](#). 2019;17(1):34-40.
51. Yeşilyurt AM, Pehlivan N, Durmuş N, KARAOĞLU SA. Trichoderma citrinoviride: a potent biopriming agent for the alleviation of salt stress in maize. [Hacettepe Journal of Biology and Chemistry](#). 2018 Jan 3;46(1):101-11.
52. Bakhshandeh E, Gholamhosseini M, Yaghoobian Y, Pirdashti H. Plant growth promoting microorganisms can improve germination, seedling growth and potassium uptake of soybean under drought and salt stress. [Plant Growth Regulation](#). 2020 Jan;90:123-36.
53. Meena M, Swapnil P, Divyanshu K, Kumar S, Harish, Tripathi YN, Zehra A, Marwal A, Upadhyay RS. PGPR-mediated induction of systemic resistance and physiochemical alterations in plants against the pathogens: Current perspectives. [Journal of Basic Microbiology](#). 2020 Oct;60(10):828-61.
54. Galeano RM, Franco DG, Chaves PO, Giannesi GC, Masui DC, Ruller R, Corrêa BO, da Silva Brasil M, Zanoelo FF. Plant growth promoting potential of endophytic *Aspergillus niger* 9-p isolated from native forage grass in Pantanal of Nhecolândia region, Brazil. [Rhizosphere](#). 2021 Jun 1;18:100332.

How to cite this article

Z. Rezaloo, S. Shahbazi, A. A. Aliloo, M. Sarajuoghi, *Impact of Mutants of Trichoderma Species in Modulating Salinity Stress of Beans (Phaseolus vulgaris L. CV. Khomein)*, Journal of Nuclear Science and Applications (JONRA), Vol. 3, No. 4, P 47-58, Autumn (2023),
Url: https://jonra.nstri.ir/article_1546.html, DOI: <https://doi.org/10.24200/jon.2023.1075>.



This work is licensed under the Creative Commons Attribution 4.0 International License.
To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0>